



Corporate Headquarters

5400 Kennedy Avenue
Cincinnati, Ohio 45213

Phone: 513-281-3400
Toll-Free: 1-877-PROSCAN (776-7226)
Fax: 513-281-3420
Web: www.proscan.com

Imaging Sites

Ohio

Central Ohio

- Dublin
- Gahanna
- Pickerington
- Polaris

Cincinnati

- Eastgate
- Midtown
- Mason
- Paul Brown Stadium
- Tri-County
- Tyersville
- Western Hills

Troy

Arkansas

- North Little Rock

Florida

- Bonita Springs
- Naples North
- Naples South

Indiana

Indianapolis

- Avon
- Carmel
- Fishers
- Greenwood

Kentucky

- Lexington

Louisiana

Maryland

- Columbia

New York

- Buffalo

To Our Breast Feeding patients,

After careful study of the Manual on Contrast Media provided by the American College of Radiology (ACR) we follow the recommendations of the ACR. The findings show that a very small amount of all Iodinated contrast media can be excreted in breast milk up to 24 hours after injection and absorbed by the infant's gut. According to the ACR guidelines, it is safe for the mother and infant to continue breast feeding after receiving such an agent. Ultimately, an informed decision to temporarily stop breast feeding should be left up to the mother after all facts are communicated. Our hope is that by advising you of this protocol in advance of your procedure you may make the choice that is best for you and your child and will have an opportunity to express and store breast milk for use after the examination if so desired. Excerpts of the guidelines are provided for your review as well as the link to the ACR website if you want to read further on the issue.

Respectfully,
PROSCAN IMAGING



The Most Trusted Name In Medical Imaging



<https://www.acr.org/Quality-Safety/Resources/Contrast-Manual>

Iodinated X-ray Contrast Media

Background

The plasma half-life of intravenously administered iodinated contrast medium is approximately 2 hours, with nearly 100% of the media cleared from the bloodstream in patients with normal renal function within 24 hours. Because of its low lipid solubility, less than 1% of the administered maternal dose of iodinated contrast medium is excreted into the breast milk in the first 24 hours. In addition, less than 1% of the contrast medium ingested by the infant is absorbed from its gastrointestinal tract. Therefore, the expected systemic dose absorbed by the infant from the breast milk is less than 0.01% of the intravascular dose given to the mother. This amount represents less than 1% of the recommended dose for an infant being prescribed iodinated contrast material related to an imaging study (usually 1.5 to 2 mL/kg). The potential risks to the infant include direct toxicity and allergic sensitization or reaction, which are theoretical concerns but have not been reported.

The likelihood of either direct toxic or allergic-like manifestations resulting from ingested iodinated contrast material in the infant is extremely low. As with other medications in milk, the taste of the milk may be altered if it contains contrast medium.

Recommendation

Because of the very small percentage of iodinated contrast medium that is excreted into the breast milk and absorbed by the infant's gut, we believe that the available data suggest that it is safe for the mother and infant to continue breast-feeding after receiving such an agent.

Ultimately, an informed decision to temporarily stop breast-feeding should be left up to the mother after these facts are communicated. If the mother remains concerned about any potential ill effects to the infant, she may abstain from breast-feeding from the time of contrast administration for a period of 12 to 24 hours. There is no value to stop breast feeding beyond 24 hours. The mother should be told to express and discard breast milk from both breasts during that period. In anticipation of this, she may wish to use a breast pump to obtain milk before the contrast-enhanced study to feed the infant during the 24-hour period following the examination.